

Policy on Model Dental Benefits for Infants, Children, Adolescents, and Individuals With Special Health Care Needs

Originating Councils

Council on Dental Benefit Programs/Council on Clinical Affairs

Adopted

2008

Purpose

The American Academy of Pediatric Dentistry (AAPD) believes that all infants, children, adolescents, and individuals with special health care needs must have access to comprehensive preventive and therapeutic oral health care benefits that contribute to their optimal health and well-being. This policy is intended to assist policy makers, third-party payors, and consumer groups/benefits purchasers to make informed decisions about the appropriateness of oral health care services for these patient populations.

Methods

This policy is based upon a review of AAPD's systematically-developed oral health policies and clinical practice guidelines as well as clinical practice guidelines that have been developed by other professional organizations and endorsed by the AAPD.

Background

The AAPD, in accordance with its vision and mission, advocates optimal oral health and health care for all infants, children, adolescents, and individuals with special health care needs. Oral diseases are progressive and cumulative; ignoring oral health problems can lead to needless pain and suffering, infection, loss of function, increased health care costs, and life-long consequences in educational, social, and occupational environments. When oral health care is not accessible, the health implications, effects on quality of life, and societal costs are enormous.¹ The AAPD's oral health policies and clinical guidelines² encourage the highest possible level of care to children and patients with special health care needs. The AAPD also sponsors a national consensus conference or symposium each year on pediatric oral health care and publishes those proceedings in a special issue of *Pediatric Dentistry*. Those documents,²⁻⁶ as well as clinical practice guidelines from other organizations with recognized professional expertise and stature,⁷⁻¹³ serve as the basis for the recommendations below. Such recommendations ideally are evidence based but, in the absence of conclusive evidence, may rely on expert opinion and clinical observations.

Policy Statement

The AAPD encourages all policy makers and third party payors to consult the AAPD in the development of benefit plans that best

serve the oral health interests of infants, children, adolescents, and individuals with special health care needs. These model services are predicated on establishment of a dental home, defined as the ongoing relationship between the dentist (ie, the primary oral health care provider) and the patient, inclusive of all aspects of oral health care, starting no later than 12 months of age.¹⁴

Expected benefits of care should outweigh potential risks. Value of services is an important consideration, and all stakeholders should recognize that cost-effective care is not necessarily the least expensive treatment. Consistent with AAPD clinical guidelines, the following services should be included in health benefit plans.

A. Preventive services:

- (1) Initial and periodic examinations of the dentition and oral cavity, including medical and dental histories, furnished in accordance with the attached periodicity schedule¹⁵ or when oral screenings by other health care providers indicate a risk of caries or other dental or oral disease;
- (2) Education for the patient and the patient's family on measures that promote oral health as part of initial and periodic well-child assessment;
- (3) Age-appropriate anticipatory guidance and counseling on non-nutritive habits, injury prevention, and tobacco use/substance abuse;
- (4) Application of topical fluoride at a frequency based upon caries risk factors;
- (5) Prescription of dietary fluoride supplement¹² based upon a child's age, caries risk, and fluoride level of the water supply or supplies;
- (6) Application of pit and fissure sealants based on caries risk factors, not based upon patient age or time lapsed since eruption;¹⁶
- (7) Dental prophylactic services at a frequency based on caries and periodontal risk factors.

B. Diagnostic procedures consistent with guidelines developed by organizations with recognized professional expertise and stature, including radiographs in accordance with recommendations by the US Food and Drug Administration and the American Dental Association.⁹

C. Restorative and endodontic services to relieve pain, resolve infection, restore teeth, and maintain dental function and oral health. This would include interim therapeutic restorations, a beneficial provisional technique in contemporary pediatric restorative dentistry.¹⁷

D. Orthodontic services including space maintenance and services to diagnose, prevent, intercept, and treat malocclusions, including management of children with cleft lip or palate and/or congenital or developmental defects. These services include, but are not limited to, initial appliance construction and replacement of appliances as the child grows.

E. Dental and oral surgery which shall include sedation/general anesthesia and related medical services that shall be furnished on an inpatient basis when medically necessary.

F. Periodontal services to resolve gingivitis, periodontitis, and other periodontal diseases or conditions in children.

G. Prosthodontic services, including implants to restore oral function, that are consistent with guidelines developed by organizations with recognized professional expertise and stature.

H. Diagnostic and therapeutic services related to the management of orofacial trauma. When the injury involves a primary tooth, benefits should cover complications for the developing succedaneous tooth.

I. Drug prescription for preventive services, relief of pain, or treatment of infection.

J. Medically necessary services for preventive and therapeutic care in patients with medical, physical, or behavioral conditions. These services include, but are not limited to, the care of hospitalized patients, sedation, and general anesthesia in outpatient or inpatient hospital facilities.

K. Behavior guidance services necessary for the provision of optimal therapeutic and preventive oral care to patients with medical, physical, or behavioral conditions. These services may include both pharmacologic and non-pharmacologic management techniques.

L. Consultative services provided by a pediatric dentist when the dental home has been established with a general practitioner or when requested by another dental specialist or medical care provider.

References

1. US Dept of Health and Human Services. Oral health in America: A report of the Surgeon General—Executive summary. Rockville, Md: US Dept of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000. Available at: "<http://www2.nidcr.nih.gov/sgr/execsumm.htm>". Accessed April 3, 2008.
2. American Academy of Pediatric Dentistry. Reference manual. Pediatric Dentistry Supplement. Chicago, IL. Available at: "<http://www.aapd.org/media/policies.asp>". Accessed June 7, 2008.
3. American Academy of Pediatric Dentistry. Pediatric dentistry restorative consensus conference. *Pediatr Dent* 2002; 24(5):374-516.
4. American Academy of Pediatric Dentistry. Symposium on behavior guidance. *Pediatr Dent* 2004;26(2):110-83.
5. American Academy of Pediatric Dentistry. Symposium on the prevention of oral disease in children and adolescents. *Pediatr Dent* 2006;28(2):96-198.
6. American Academy of Pediatric Dentistry, American Association of Endodontists. Proceedings of the Joint Symposium on Emerging Science in Pulp Therapy: New Insights into Dilemmas and Controversies. *Pediatr Dent* 2008;30(3): 192-267.
7. American Association of Endodontists. Guide to Clinical Endodontics. 4th ed. Chicago, Ill. American Assoc of Endodontists 2004.
8. American Academy of Periodontology. Periodontal diseases of children and adolescents. *J Periodontol* 2003;74:1696-704.
9. American Dental Association, US Dept of Health and Human Services. The Selection of Patients for X-Ray Examination: Dental Radiographic Examinations. Rockville, Md: Food and Drug Administration, 2004; HHS Publication Number 88-8273. Available at: "<http://www.ada.org/prof/resources/topics/radiography.asp>". Accessed August 8, 2008.
10. American Cleft Palate-Craniofacial Association. Parameters for evaluation and treatment of patients with cleft lip/palate or other craniofacial anomalies. Chapel Hill, NC: The Maternal and Child Health Bureau, Title V, Social Security Act, Health Resources and Services Administration, US Public Health Service, DHHS; November 2007. Grant # MCJ-425074. Available at: "www.acpa-cpf.org/teamcare/parameters07rev.pdf". Accessed April 1, 2008.
11. National Foundation for Ectodermal Dysplasias. Parameters of oral health care for individuals affected by ectodermal dysplasias. National Foundation for Ectodermal Dysplasias. Mascoutah, Ill. 2003. Available at: "www.nfed.org". Accessed June 7, 2008.
12. American Academy of Pediatrics Committee on Nutrition. Fluoride supplementation for children: Interim policy recommendations. *Pediatrics* 1995;95(5):777.
13. Casamassimo P, Holt K, eds. Bright Futures in Practice: Oral Health—Pocket Guide. Washington, DC: National Maternal and Child Oral Health Resource Center 2004.

- 14. American Academy of Pediatric Dentistry. Policy on the dental home. *Pediatr Dent* 2008;30(suppl):22-3.
- 15. American Academy of Pediatric Dentistry. Guideline on periodicity of examination, preventive dental services, anticipatory guidance, and oral treatment for infants, children, and adolescents. *Pediatr Dent* 2007;29(suppl):102-8.
- 16. Feigal RJ, Donly KJ. The use of pit and fissure sealants. *Pediatr Dent* 2006;28(2):143-50.
- 17. American Academy of Pediatric Dentistry. Guideline on pediatric restorative dentistry. *Pediatr Dent* 2008;30(suppl):163-9.

RECOMMENDATIONS FOR PREVENTIVE PEDIATRIC ORAL HEALTH CARE

Since each child is unique, these recommendations are designed for the care of children who have no contributing medical conditions and are developing normally. These recommendations will need to be modified for children with special health care needs or if disease or trauma manifests variations from normal. The American Academy of Pediatric Dentistry (AAPD) emphasizes the importance of very early professional intervention and the continuity of care based on the individualized needs of the child.

	AGE	6–12 MONTHS	12–24 MONTHS	2–6 YEARS	6–12 YEARS	12 YEARS AND OLDER
Clinical oral examination ^{1,2}		o	o	o	o	o
Assess oral growth and development ³		o	o	o	o	o
Caries-risk assessment ⁴		o	o	o	o	o
Radiographic assessment ⁵		o	o	o	o	o
Prophylaxis and topical fluoride ^{4,5}		o	o	o	o	o
Fluoride supplementation ^{6,7}		o	o	o	o	o
Anticipatory guidance/ counseling ⁸		o	o	o	o	o
Oral hygiene counseling ⁹		Parent	Parent	Patient/parent	Patient/parent	Patient
Dietary counseling ¹⁰		o	o	o	o	o
Injury prevention counseling ¹¹		o	o	o	o	o
Counseling for nonnutritive habits ¹²		o	o	o	o	o
Counseling for speech/language development		o	o	o		
Substance abuse counseling					o	o
Counseling for intraoral/perioral piercing					o	o
Assessment and treatment of developing malocclusion				o	o	o
Assessment for pit and fissure sealants ¹³				o	o	o
Assessment and/or removal of third molars						o
Transition to adult dental care						o

¹ First examination at the eruption of the first tooth and no later than 12 months. Repeat every 6 months or as indicated by child's risk status/susceptibility to disease.

² Includes assessment of pathology and injuries.

³ By clinical examination.

⁴ Must be repeated regularly and frequently to maximize effectiveness.

⁵ Timing, selection, and frequency determined by child's history, clinical findings, and susceptibility to oral disease.

⁶ Consider when systemic fluoride exposure is suboptimal.

⁷ Up to at least 16 years.

⁸ Appropriate discussion and counseling should be an integral part of each visit for care.

⁹ Initially, responsibility of parent; as child develops, jointly with parent; then, when indicated, only child.

¹⁰ At every appointment; initially discuss appropriate feeding practices, then the role of refined carbohydrates and frequency of snacking in caries development and childhood obesity.

¹¹ Initially play objects, pacifiers, car seats; then when learning to walk, sports and routine playing, including the importance of mouthguards.

¹² At first, discuss the need for additional sucking; digits vs pacifiers; then the need to wean from the habit before malocclusion or skeletal dysplasia occurs. For school-aged children and adolescent patients, counsel regarding any existing habits such as fingernail biting, clenching, or bruxism.

¹³ For caries-susceptible primary molars, permanent molars, premolars, and anterior teeth with deep pits and fissures; placed as soon as possible after eruption.